

The fundamental group of the harmonic archipelago

Paul Fabel

Department of Mathematics & Statistics
Mississippi State University

Abstract

The harmonic archipelago HA is obtained by attaching a large pinched annulus to every pair of consecutive loops of the Hawaiian earring. We clarify $\pi_1(HA)$ as a quotient of the Hawaiian earring group, provide a precise description of the kernel, show that both $\pi_1(HA)$ and the kernel are uncountable, and that $\pi_1(HA)$ has the indiscrete topology.

1 Introduction

This note serves to clarify certain properties of $\pi_1(HA)$ and $\pi_1(HE)$, the topological fundamental groups respectively of the harmonic archipelago and the Hawaiian earring.

The Hawaiian earring HE is the union of a null sequence of simple closed curves meeting in a common point.

Introduced by Bogley and Sieradski, the harmonic archipelago HA , is the space obtained by attaching large pinched annuli, one for each pair of consecutive loops, to the Hawaiian earring HE . In [2] Bogley and Sieradski construct a comprehensive theory which provides a useful framework for investigating the fundamental groups of locally complicated spaces such as HE and HA . Various properties of $\pi_1(HA)$, such as its uncountability, are uncovered in [2].

In [1], Biss also uses HE and HA as motivating examples, and proves some nice general results on topological fundamental groups and their generalized covering spaces.

However, an oversight in [1] (see Remark 2), leads to a false description of $\pi_1(HA)$ and its false generalization Theorem 8.1. There is also a typographical error in the description of $\ker(j^*)$, the kernel of the epimorphism $j^* : \pi_1(HE) \rightarrow \pi_1(HA)$, induced by inclusion $j : HE \hookrightarrow HA$.

After adjusting for this, there remains arguably room for further discussion regarding which elements of $\pi_1(HE)$ belong to $\ker(j^*)$.

For example it follows from the investigations of Morgan/Morrison [6], DeSmit [4], and Cannon/Conner [3] that elements of $\pi_1(HE)$ can be seen as “transfinite words” over an infinite alphabet $\{x_1, x_2, \dots\}$ with each letter appearing finitely many times.

As described in [1], $\ker(j^*)$ is generated by the relations $x_i = x_j$ for all i and j . Does this mean two transfinite words over $\{x_1, x_2, \dots\}$ are equivalent in $\pi_1(HA)$ if and only if one can be transformed into the other by finitely many substitutions and finitely many cancellations of consecutive letters? No, for example $x_1x_2^{-1}(x_3x_4\dots)x_1x_2^{-1}(\dots x_4^{-1}x_3^{-1})$ is trivial in $\pi_1(HA)$, but cannot be transformed into the trivial word with finitely many such operations.

Are two transfinite words over $\{x_1, x_2, \dots\}$ equivalent in $\pi_1(HA)$ if one can be transformed into the other after “infinitely many substitutions”? No, for then the essential element $(x_1x_2^{-1}x_3x_4^{-1}\dots)$ could be transformed into the inessential $x_1x_1^{-1}x_2x_2^{-1}\dots$.

We provide a precise description of $\ker(j^*)$ in Theorem 7 and prove as Corollary 8 that $\ker(j^*)$ is uncountable. Corollaries 9 and 10 provide proofs of results also indicated in [2] and [1]: $\pi_1(HA)$ is uncountable and, despite its large cardinality, $\pi_1(HA)$ has the indiscrete topology.

2 Definitions

For $n \in \{1, 2, 3, \dots\}$ let $X_n \subset R^2$ denote the circle of radius $\frac{1}{n}$ centered at $(\frac{1}{n}, 0)$. Let $Y_n = \cup_{i=n}^{\infty} X_i$. Thus Y_n is the Hawaiian earring determined by the loops $X_n, X_{n+1}, \dots$

Let $A_n \subset R^2$ denote the closed pinched annulus bounded by $X_n \cup X_{n+1}$. Let $Y^n = Y_1 \cup A_1 \cup \dots \cup A_n$. Endow Y^n with the subspace topology inherited from R^2 .

For the **underlying set** let $HA = \cup_{n=1}^{\infty} Y^n$. However we define the topology of HA such that Y^n inherits the usual topology but such that 1) There exists a sequence $z_n \in \text{int}(A_n)$ such that $\{z_1, z_2, \dots\}$ has no subsequential limit and 2) If $p \in HA$ is a subsequential limit of the sequence $y_1, y_2, \dots$ and if $y_n \in \text{int}(A_n)$ for all n , then $p = (0, 0)$.

Let $G_n = \pi_1(Y_n, (0, 0))$.

Let $F_{N,n}$ denote the free group on the letters $\{x_N, x_{N+1}, \dots, x_n\}$ coupled with the symbol 1 denoting the trivial element.

Let $\phi_{N,n} : F_{N,n+1} \rightarrow F_{N,n}$ denote the homomorphism such that $\phi_{N,n}(x_i) = x_i$ if $N \leq i \leq n$ and $\phi_n(x_{n+1}) = 1$.

For $N \geq 1$ let G^N denote the inverse limit of free groups determined by $F_{N,N} \leftarrow F_{N,N+1} \leftarrow \dots$ under the bonding maps $\phi_{N,n}$.

Formally elements of $F_{N,n}$ are equivalence classes of words under

the obvious cancellations, and the group operation is concatenation. However each element of $F_{N,n}$ has a unique representative with a minimal number of nontrivial letters. Consequently each element of G^N is uniquely determined by a **canonical sequence** $w_N, w_{N+1}, \dots$ such that $w_n \in F_{N,n}$ and w_n is a maximally reduced word in $F_{N,n}$.

3 $\pi_1(HA)$

It is a nontrivial fact ([3], [4] [6]) that $\pi_1(Y_1)$ injects naturally into the inverse limit of free groups.

Remark 1 Given $n \geq N \geq 1$ let $Z_{N,n} = \cup_{i=N}^n X_n$. Let $r_{N,n} : Y_N \rightarrow Z_{N,n}$ denote the retraction collapsing X_i to $(0,0)$ for $i > n$. Since $r_{N,n}(r_{N,n+1}) = r_{N,n}$, the maps $r_{N,n}$ induce a homomorphism $\psi_N : \pi_1(Y_N, (0,0)) \rightarrow \lim_{\leftarrow} \pi_1(Z_{N,n}, (0,0))$. It is shown in [6] and [4] that ψ_N is one to one. Moreover $G_N = \text{im}(\psi_N)$ consists of all canonical sequences $(w_N, w_{N+1}, \dots)$ such that for each i there exists M_i such that for all $n \geq N$, x_i appears at most M_i times in w_n .

It is falsely asserted in [1] that $\psi_1 : \pi_1(HE) \rightarrow \lim_{\leftarrow} F_{1,n}$ is an isomorphism. Consequently Theorem 8.1 of [1] is false.

Remark 2 The canonical monomorphism $\psi_1 : \pi_1(Y_1, (0,0)) \hookrightarrow G^1$ is **not** surjective. By compactness of $[0, 1]$, a given path in Y_1 can traverse each loop only finitely many times. Thus the element

$$(1, x_1 x_2 x_1^{-1} x_2^{-1}, x_1 x_2 x_1^{-1} x_2^{-1} x_1 x_3 x_1^{-1} x_3^{-1}, \dots) \in \lim_{\leftarrow} F_{1,n}$$

has no preimage in the Hawaiian earring group.

Remark 3 $X_n \cup X_{n+1}$ is a strong deformation retract of $A_n \setminus \{z_n\}$. Thus Y^n is a strong deformation retract of $HA \setminus \{z_{n+1}, z_{n+2}, \dots\}$.

Lemma 4 Suppose $f : S^1 \rightarrow Y_1$ is any map. Then f is inessential in HA if and only if there exists N such that f is inessential in Y^N .

Proof. Suppose f is inessential in HA . Let $F : D^2 \rightarrow HA$ be a continuous extension of f . Since $\text{im}(F)$ is compact, there exists N such that $z_n \notin \text{im}(F)$ whenever $n > N$. Thus $\text{im}(F) \subset HA \setminus \{z_{N+1}, z_{N+2}, \dots\}$. By remark 3 Y^N is a strong deformation retract of $HA \setminus \{z_{N+1}, z_{N+2}, \dots\}$. Thus f is inessential in Y^N . Conversely if there exists N such that f is inessential in Y^N then f is inessential in HA since $Y^N \subset HA$. ■

Notice Y_N is a strong deformation retract of Y^{N-1} under a homotopy $R_{N,t} : Y^{N-1} \rightarrow Y_N$ collapsing $A_1 \cup \dots \cup A_{N-1}$ onto the simple closed curve

X_N . In particular given any loop $f : S^1 \rightarrow Y_1$ we may canonically deform f in Y^{N-1} to a loop g such that $\text{im}(g) \subset Y_N$. Appealing to Remark 1 we may identify $\pi_1(Y_i, (0, 0))$ with G_i . Thus the composition $Y_1 \hookrightarrow Y^{N-1} \rightarrow Y_N$ induces a homomorphism $q_N^* : G_1 \rightarrow G_N$. Combining these observations we obtain the following:

Lemma 5 $q_N^*(w_1, w_2, \dots) = (v_N, v_{N+1}, \dots)$ if and only if the following property is satisfied for each $n \geq N$: For each $i \leq N$ replace each occurrence of x_i in w_n with x_N , creating a word $\hat{w}_n$ on the letters $\{x_N, \dots, x_n\}$. Then the word $\hat{w}_n$ is equivalent to v_n in the free group $F_{N,n}$.

Lemma 6 is also handled in [1] and [2]. Our proof is similar to an argument that the 2 sphere is simply connected.

Lemma 6 Let $j : Y_1 \hookrightarrow HA$ denote the inclusion map. Then the induced homomorphism $j^* : \pi_1(Y_1, (0, 0)) \rightarrow \pi_1(HA, (0, 0))$ is surjective.

Proof. Suppose $f : S^1 \rightarrow HA$ is any map such that $f(1) = (0, 0)$. Let J be a (nonempty) component of $f^{-1}(HA \setminus Y_1)$. Note $HA \setminus Y_1$ is the union of pairwise disjoint connected open sets $\text{int}(A_1) \cup \text{int}(A_2), \dots$. Thus there exists i such that $f(J) \subset \text{int}(A_i)$ and $f(\partial J) \subset \partial A_i$. If $z_i \in \text{im}(f)$ replace $f|_J$ by a path homotopic small perturbation $\hat{f}_J$ such that $z_i \notin \text{im}(\hat{f}_J)$. By uniform continuity of f finitely many such surgeries are required. Note f and $\hat{f}$ are homotopic in HA and $\text{im}(\hat{f}) \subset HA \setminus \{z_1, z_2, \dots\}$.

By remark 3 Y_1 is a strong deformation retract of $HA \setminus \{z_1, z_2, \dots\}$ and hence there exists $f^* : S^1 \rightarrow Y_1$ such that $j^*[f^*] = [\hat{f}] = [f]$. ■

Since j^* is a surjection $\pi_1(HA, (0, 0))$ is isomorphic to the quotient group $\pi_1(Y_1, (0, 0)) / \ker j^*$. Combining Lemmas 4 and 5 we obtain the following characterization of $\ker j^*$.

Theorem 7 Let $j^* : G_1 \rightarrow \pi_1(HA, (0, 0))$ denote the epimorphism induced by inclusion $j : Y_1 \hookrightarrow HA$. Let $K = \ker j^*$. Then $(w_1, w_2, \dots) \in K$ if and only if there exists N such that the following holds: Suppose for each $i \leq N$ and for each $n \geq N$ each occurrence of x_i in w_n is replaced by x_N creating a (nonreduced) word v_n on the letters $\{x_N, x_{N+1}, \dots, x_n\}$. Then v_n can be reduced to the trivial element of $F_{N,n}$.

Corollary 8 $\ker(j^*)$ is uncountable.

Proof. There exist uncountably many distinct permutations of the set $\{3, 4, 5, 6, \dots\}$. Moreover, the loops in Y_3 determined by distinct permutations of $x_3, x_4, \dots$ determine distinct elements of $\pi_1(Y_3, (0, 0))$. Thus

G_3 is uncountable. Each $w \in G_3$ determines a homotopically distinct loop in Y_1 corresponding to $x_1 x_2^{-1} w x_1 x_2^{-1} w^{-1}$. By Theorem 7 this element is in $\ker(j^*)$. ■

Corollary 9 is also treated in [2].

Corollary 9 $\pi_1(HA, (0, 0))$ is uncountable.

Proof. Consider the set A of functions from $\{1, 3, 5, \dots\} \rightarrow \{0, 1\}$. Each element $f \in A$ determines a permutation $\tau_f : \{1, 2, 3, \dots\} \rightarrow \{1, 2, 3, \dots\}$ as follows: If $f(2n - 1) = 0$ then τ_f fixes $2n - 1$ and $2n$. Otherwise τ_f swaps $2n - 1$ and $2n$. The permutation τ_f determines a loop in Y_1 by the following recipe. Travel clockwise once around $X_{\tau_f(1)}$, then travel clockwise once around $X_{\tau_f(2)}$, etc... This corresponds to the transfinite word $\tau_f(1)\tau_f(2)\dots \in \pi_1(Y_1)$.

Note A is uncountable. Declare two elements $\{f, g\} \subset A$ equivalent if f and g agree except on a finite set. Each equivalence class in A has countably many elements and hence B , the set of equivalence classes of A , is uncountable. If $[f]$ and $[g]$ are distinct elements of B then by Lemma 5, for each N , τ_f and τ_g fail to be equivalent in Y^N . Thus by Lemma 4 τ_f and τ_g determine distinct elements of $\pi_1(HA, (0, 0))$. Thus, since B is uncountable, $\pi_1(HA, (0, 0))$ is uncountable. ■

It shown in [1] (and generalized in [5]) that the quotient space consisting of path components of based loops in a topological space X is a topological group. Despite a large number of elements, $\pi_1(HA, (0, 0))$ has the indiscrete topology. Corollary 10 is also argued in [1].

Corollary 10 The topological group $\pi_1(HA, (0, 0))$ has exactly one nonempty open subset.

Proof. Let $L = \{f : [0, 1] \rightarrow HA \mid f(0) = f(1) = (0, 0)\}$ with the uniform topology. By definition $\pi_1(HA, (0, 0))$ is the collection of path components of L with the quotient topology. Suppose $B \subset \pi_1(HA, (0, 0))$, suppose $B \neq \emptyset$ and suppose $[f] \in B$.

Since $im(f)$ is compact and $\{z_1, z_2, \dots\}$ is not compact, there exists N such that $im(f) \subset HA \setminus \{z_{N+1}, z_{N+2}, \dots\}$. For $M > N$ there is a strong deformation retraction from $HA \setminus \{z_{M+1}, z_{M+2}, \dots\}$ onto Y^M . Thus for large M we may deform f in HA to a nearby map f_M such that $im(f_M) \subset Y^M$. Let $A_M = f_M^{-1}(Y^M \setminus (X_{M+1} \cup X_{M+2} \dots))$. Redefine f_M over A_M to be the constant $(0, 0)$ creating a map $\hat{f}_M$. Note $im(\hat{f}_M) \subset A_1 \cup \dots \cup A_{M-1}$. Notice $A_1 \cup \dots \cup A_{M-1}$ has the homotopy type of S^1 . By wrapping around X_M as many times as necessary, we may extend $\hat{f}_M$ to a map of $f_M^* : [0, 1 + \frac{1}{M}] \rightarrow HA$ such that f_M^* is an inessential loop and $f(t) \in X_M$ for $t \in [1, \frac{1}{M}]$. Reparameterize f_M^* linearly to create a

map $f_M^{**} : [0, 1] \rightarrow HA$. Notice $f_M^{**} \rightarrow f$ uniformly. Thus the path component of the trivial loop is dense in the space of based loops of HA . Thus, if B is open then B contains the trivial element.

Suppose on the other hand that B is closed. Choose N such that $im(f) \subset Y^n$ whenever $n \geq N$. Notice Y_{n+1} is a strong deformation retract of Y^n . Thus $[f]$ contains representatives whose images have arbitrarily small diameter. Thus B contains the trivial element.

Thus the trivial element of $\pi_1(HA(0, 0))$ belongs to each nonempty open set and each nonempty closed set. Hence $\pi_1(HA(0, 0))$ has only one nonempty open subset. ■

References

- [1] Biss, Daniel K. The topological fundamental group and generalized covering spaces. *Topology Appl.* 124 (2002), no. 3, 355–371.
- [2] Bogley, W.A., Sieradski, A.J. Universal path spaces. Preprint. <http://oregonstate.edu/~bogleyw/#research>.
- [3] Cannon, J. W.; Conner, G. R. The combinatorial structure of the Hawaiian earring group. *Topology Appl.* 106 (2000), no. 3, 225–271.
- [4] de Smit, Bart. The fundamental group of the Hawaiian earring is not free. *Internat. J. Algebra Comput.* 2 (1992), no. 1, 33–37.
- [5] Fabel, Paul. Completing Artin’s braid group on infinitely many strands. <http://front.math.ucdavis.edu/math.GT/0201303>. To appear in *Journal of Knot Theory and its Ramifications*.
- [6] Morgan, John W.; Morrison, Ian. A van Kampen theorem for weak joins. *Proc. London Math. Soc.* (3) 53 (1986), no. 3, 562–576.